

## Further Use of Natural Infrasound as a Continuous Monitor of the Upper Atmosphere<sup>1,2</sup>

DAVID RIND

*Lamont-Doherty Geological Observatory of Columbia University, Palisades, N. Y. 10964*

WILLIAM L. DONN

*Lamont-Doherty Geological Observatory of Columbia University and  
The City College of the City University of New York, New York, N. Y. 10031*

(Manuscript received 13 January 1975, in revised form 16 April 1975)

### ABSTRACT

This is a further study of the use of natural infrasound in the atmosphere to monitor tidal circulation in the lower thermosphere. The height of this circulation is determined with the use of a reference atmospheric model, which is then used to calibrate infrasound/microseism ratios in terms of height. Also, we show from continuous observation over six years at 41°N, 74°W that the winter semidiurnal tide is present at least 62% of the time and the diurnal, at least 42% of the time.

### 1. Introduction

This paper reports the attempts to refine the infrasound technique for monitoring upper atmosphere winds and temperatures (Donn and Rind, 1972; Rind *et al.*, 1973). Naturally occurring infrasound is generated continuously by ocean waves and propagates up to the stratosphere or lower thermosphere where it undergoes reflection from the combined effects of wind and temperature. In this paper we will discuss several methods for estimating the level of reflection. We then apply these results to determine the regularity of upper air circulations, which is possible because the system is in operation 24 h a day, every day of the year.

Infrasound has been recorded for the past eight years at Palisades, N. Y. (41°N, 74°W), with the use of capacitor microphones arranged in a tripartite array. Relevant details are available in Donn and Rind (1972). Cross correlation of the data recorded both on magnetic tape and on visible chart records allows the direction and trace velocity of the signal to be determined. We have already shown (Rind *et al.*, 1973) that the trace velocity is equal to the speed of sound which is a function of wind and temperature at the reflection level. In all of this work, a crucial problem is the determination of the level of reflection; this is discussed in detail below.

### 2. Absolute height determination

Reflection of infrasound originating at the surface occurs when the sound velocity at any level exceeds the sound velocity at the surface. The sound velocity in any direction  $V(\mathbf{i})$  is equal to the component of the wind velocity  $W$  in that direction plus the sound speed  $C$ . Thus,

$$V(\mathbf{i}) = W\mathbf{i} + C, \quad (1)$$

where  $C = (\gamma RT/m)^{1/2}$ ,  $T$  is the absolute temperature,  $R$  the universal gas constant,  $\gamma = c_p/c_v$ , and  $m$  is the mean molecular weight of air. To determine the probable level of reflection we must know the distribution of these parameters in the vertical. Thus, we are initially dependent on observations collected by other methods. For the purpose of composing such a model we conducted an extensive survey of published wind and temperature observations; the relevant sources, techniques and locations are given in Table A1 in the Appendix. The model was composed for our latitude and longitude for each month of the year; it includes prevailing effects and semidiurnal and diurnal tides for both wind and temperature at 5 km height intervals between 0 and 120 km. We used data for other latitudes bearing in mind the expected latitudinal variation of the particular phenomenon (for instance the  $S_{2,4}$  semidiurnal tide). The ionospheric drift observations were used with due consideration for all the ambiguities involved in their interpretations. Finally, we compared our model with the prevailing wind models of Groves (1971) and Theon (1972) [and others listed in Table A] and

<sup>1</sup>Lamont-Doherty Geological Observatory of Columbia University Contribution No. 2233.

<sup>2</sup>University Institute of Oceanography, City University of New York, Contribution No. 58.




FIG. 1. Sound velocity vs height for February (solid line) and July (dotted line) computed from wind and temperature model.

mean temperature models of Champion (1967), Cole and Kantor (1972) and Salah *et al.* (1974), among others, to attempt to clarify any discrepancies.

Fig. 1 shows the model results for sound velocity from the east (where our strongest sources are located) for prevailing conditions calculated with the use of (1) for February and July (February rather than January is used to reduce the influence of stratospheric warmings in this discussion).

In summer, because the average sound velocity is greater than that at ground level from about 38 to 65 km, it is within this region that the signal will be reflected. In winter, because of the strong stratospheric west winds below, reflection first becomes possible only above 100 km. As the temperature rises strongly in the lower thermosphere, reflection should




FIG. 2. Dissipation vs height for signal of 4 s period for two reflections calculated from standard dissipation coefficients (*U. S. Standard Atmosphere Supplements 1966*).

TABLE 1. Model of semidiurnal tide.

	Elevation (km)							
	85	90	95	100	105	110	115	120
Time of maximum west wind	1100	0930	0800	0630	0430	0300	0200	0100
Time of maximum east wind	{ 1700 0500	1530 0330	1400 0200	1230 0030	1030 2230	0900 2100	0800 2000	0700 1900
Velocity ( $\text{m s}^{-1}$ )	15	23	31	38	47	49	49	47

be certain from above about 110 km. However, this is a region of strong dissipation for signal of the frequency recorded (Donn and Rind, 1972). Fig. 2 shows the cumulative dissipation for sound of this frequency along its path through these levels. Clearly, signal recorded from reflections above 110 km should be weak; furthermore, changes in the reflection level of 5 km should be readily apparent as changes of amplitude are dependent upon the height-related dissipation that occurs.

The exact level of reflection is, of course, determined by the actual temperature and wind at any time. This depends upon tidal components also, not simply prevailing conditions. Table 1 shows the model we synthesized for the semidiurnal tide for the region of

possible reflection in winter. For comparison, one can refer to Goodwin (1968), Groves (1971), Bernard (1974a, b) and Fellows *et al.* (1974), among others. The semidiurnal tide appears to be a stable feature and there is not much disagreement about its phase at these heights. If one adds the model in Table 1 to that of the prevailing wind and temperature in Fig. 1 the following can be expected: as the tidal wind becomes east at levels close to 100 km, the signal will be reflected from these heights rather than from higher, more dissipative regions, where the high temperatures always provide a sound reflection level. The amplitude of the signal should thus show a semidiurnal variation in agreement with the tidal east wind descent evident in Table 1.




FIG. 3. Average infrasound amplitude (arbitrary units) vs time of day for February from 6 years of data.

TABLE 2. Directional rotation of infrasound.

	Hour											
	06	08	10	12	14	16	18	20	22	24	02	04
Average azimuth of signal from east	85°	92°	109°	120°	140°	115°	110°	86°	93°	107°	120°	133°
Percent signal from west	25	06	06	12	15	43	39	20	12	31	17	26

Fig. 3 shows the average February infrasound amplitude as a function of the hour of the day (in arbitrary units) from 6 years of data. It can be seen that the variation has a strong semidiurnal character with maximum amplitude at 1000-1200 and 2100-2400 local time. Comparison with the phase model in Table 1 indicates strongly that the high-amplitude signal from the east comes from heights of about 105 km.

If this effect is really due to the rotating tidal influence, it should be apparent in several ways. As the direction of the semidiurnal tidal wind rotates, the direction of the signal capable of being reflected should change very slightly for a sole narrow source but more distinctly for a broad or multi-source situation. If we average over prolonged periods, a wide scatter of sources will exist, and a directional rotation of infrasound should become evident. In Table 2 constructed from computed azimuths by averaging 345 h of data scattered over 6 years such an effect is shown. The most southerly (130°-140°) azimuth agrees with a semidiurnal meridional wind in quadrature with that shown in Table 1 for 105 km.

One final presentation of this rotational effect can be made. In Fig. 4 the amplitude variation versus time is displayed for signals from the northeast, east and southeast. The phase of the reflecting (tidal) wind responsible for such variations agrees with results shown above and fixes the average region of reflection at 105 km for the high-amplitude signal, and above this level for signal of lower amplitude. These two studies verify further that our interpretation of reflection provided by the rotating semi-diurnal tide is valid.

### 3. "Independent" height determination

Having determined the approximate height of reflection from data determined by other techniques we are now in a position to apply this result to develop an independent method. This method involves relating the signal amplitude to the source strength as indicated by microseisms generated simultaneously. Microseisms are short-period seismic waves radiated through the crust of the earth from the effect of inter-




FIG. 4. Average amplitude (arbitrary units) vs time of day for infrasound from the northeast (dotted line), east (solid line) and southeast (dashed line).

TABLE 3. Infrasound parameters for various atmospheric situations.

	Winter (120 h of data)			Summer (40 h of data)
	Maximum (1100, 2300)	Intermediate (0900, 2000)	Minimum (0700, 1700)	
Ratio ( $I/M$ )	0.4	0.2	0.1	0.54
Estimated reflection height (km)	105	110	110+	45-50
Average sound velocity ( $\text{m s}^{-1}$ )	350	348	335	365
Average direction (deg)	96	90	97	105

fering ocean waves on the sea bottom. According to Posmentier (1967) and Donn and Naini (1973), the storm waves simultaneously generate atmospheric infrasound of the same period as the microseisms.

The main parameter in the "independent" method of height determination is the ratio of infrasound to microseism amplitude ( $I/M$ ). We first determine this ratio for signals known, from the model described, to be reflected at 105 km with about a 5 km height resolution. As this ratio rises and falls we estimate the reflection level lowering and rising respectively below or above 105 km. This study is still in its infancy; the data presented here are a preliminary indication of what may be accomplished.

At Lamont, standard recording of the three components of ground motion (vertical, east-west and north-south) is made. Because microseisms contain a dominant Rayleigh wave component one can correlate the different ground motions to determine the direction of the signal. This has been done using an analogue correlator. When the direction of the microseisms was the same as that of the infrasound determined from our tripartite array (which was often the case), the amplitudes (power) were compared using the autocorrelation values. The  $I/M$  ratios determined for different times of the day are shown in Table 3 for winter. We note that the changing ratio in winter is in accord with the physical description of changing reflection height during the day in a dissipative atmosphere. The ratio for summer is greater, indicative of the lower level of reflection evident in Fig. 1. It is this type of analysis, applied on a continual basis, that allows the possibility of estimating reflection heights independently, once the "model" calibration is performed which specifies the height of reflection for a given  $I/M$  ratio.

#### 4. Atmospheric tidal circulation in winter

Having determined the approximate height of reflection we are in a position to relate some statistical summaries of circulation patterns that predominate at 100-115 km.

##### a. Semidiurnal oscillation

In our latitude ( $40^\circ\text{N}$ ) the semidiurnal tide is quite effective in causing amplitude variations of infrasound

in the manner previously described. As we record the signal continuously it is possible to give a meaningful representation of how often this tide provides for reflection from the 105 km region. Between October and mid-April the ocean is nearly always agitated enough to generate relatively strong infrasound as shown by the presence of microseisms; hence the prominent amplitude variations due to tidal effects ought to be observed. Thus, if we fail to see an amplitude oscillation with semidiurnal character this could mean one of two things: either the tide is absent or the prevailing wind and temperature have altered sufficiently to prevent tidal effects from causing reflection below the dissipation level. We cannot distinguish between these two effects but merely note the percentage of times the semidiurnal tide allows for the normal pattern of amplitude variation with east winds at 105 km about 1200 and 2400 LT. In winter (defined here as October to mid-April) analysis shows that this occurs 62% of the time. Thus the semidiurnal tide as modelled by Table 1 is a fairly regular feature of the winter circulation. Because prevailing wind and temperature changes presumably do occur, this percentage should be thought of as the lower limit of semidiurnal tide reliability.

Note that we have referred to the semidiurnal oscillation as being one of wind; incoherent scatter results (Fontanari and Alcayde, 1974; Bernard, 1974a) show that a semidiurnal oscillation in temperature also exists. Its amplitude is of the order of 20-30 K; in view of the expected tidal wind variation of  $45 \text{ m s}^{-1}$ , the square-root dependence of sound velocity on temperature indicates that the average temperature oscillation contributes only one-fourth that of the wind.

##### b. Diurnal oscillation

The existence of a diurnal oscillation in the upper winds in our vicinity is discernible in data already presented (Donn and Rind, 1972). According to Fig. 3 a different amplitude pattern is obvious in the early morning compared to the early evening; higher infrasound amplitudes at 0400 suggest east winds around 105 km. Because this is a summary of data the lower amplitude peak in Fig. 3 at 0400 compared with the two main peaks is indicative of the sporadic nature of the occurrence.

Careful scrutiny of Table 3 reveals that the directional rotation is not 12 h apart but is quicker in the morning than afternoon (120° azimuth at 0200 and 1200). Also listed in Table 3, is the percent of time that the predominant signal was from the west. This signal comes from sources in the Pacific Ocean and is quite common, being reflected by stratospheric westerlies far below any level of dissipation. It is the primary signal only when signal from the east is quite weak. According to Table 3 dominant signal from the west occurred more frequently from 1600 and 1800 LT when the east signal is weak (Fig. 3) than from 0400 and 0600. All this suggests a diurnal wind component with maximum east wind at 0400 added to the predominant semidiurnal tide.

As with the semidiurnal oscillations, continuous observation allows us to investigate the regularity of the diurnal effect. In the same context as the percentage given for the semidiurnal tide for 6 years of winter data, we see a diurnal oscillation with maximum east wind at approximately 0400 (and no corresponding effect at 1600) 42% of the time. Although the phase of this oscillation does vary, our data show that 0400-0600 is by far the most common time for maximum east winds. We conclude that the diurnal tidal wind oscillation must be of quite strong amplitude to produce east winds at 0400 in the same region that semidiurnal winds are east at 1000 and thus west at 0400.

Our observations of records during stratospheric warmings show that the diurnal tide (as well as the semidiurnal tide) has quite a strong amplitude at these times with maximum east winds invariably at 0400. This suggests that the thermospheric tide is coupled to the circulation patterns below.

**5. Summary and conclusions**

We have shown here and in previous papers that infrasound reception (from marine sources) is controlled by the behavior of the atmospheric tides.

By inversion we use infrasound behavior to determine tidal circulation. We also show how height determinations can be made on a dependent basis with reference to an atmospheric model and on an "independent" basis with the use of infrasound/microseism ratios.

The advantages of using infrasound as a monitor of the upper atmosphere are numerous. In conjunction with a technique which measures either wind or temperature, the sound velocity determination would allow the other to be known. Its continuous operation capability, inexpensive nature, and on-line data processing present the possibility of practical everyday use to determine the regularity of the diurnal and semidiurnal tide. For example, at Palisades, N. Y. (41°N, 74°W), we have concluded that the semidiurnal tide in winter is present at least 62% of the time, and the diurnal tide at least 42% of the time. These results are based on daily observations in winter for 6 years.

It would be especially interesting to compare these results with a simultaneous recording of ionospheric drifts. The close concordance of the results noted in the literature [see Briggs and Spencer (1954) for ionospheric drifts] between the two techniques convince us they look at the same level. There is no doubt that the infrasonic technique observes the neutral atmosphere winds (and temperatures) and comparison would help validate both techniques.

*Acknowledgments.* This research was supported by grants from the National Science Foundation under Contract GA 42975X and the U. S. Army Research Office-Durham under Contract DAHC-04-74-G-0172. Also, we are grateful to the Palisades Interstate Park Commission for permission to install and maintain our transducers at several sites in Tallman Mountain State Park. None of this work could have been performed without the engineering and technical services of F. Witt and B. Gallagher. Drs. N. Balachandran and A. Oppenheim read and criticized the manuscript, as well as contributing sagacious advice throughout.

APPENDIX

TABLE A1. Survey of published wind and temperature observations. (References are identified in Table A2.)

Station	Location	Technique	Reference
Carnarvon, Australia	25°N, 114°E	rocket	1
Sonmiani, Pakistan	25°N, 66°E	chaff	2
Cape Kennedy, Fla.	28°N, 80°W	rocket	3, 4, 5
Eglin AFB	29°N, 87°W	chemical trails	6-18
Hammaguir, Algeria	31°N,	chemical trails	19
Yamagawa, Japan	31°N, 131°E	ionospheric drift (I.D.)	13, 20-25
Woomera, Australia	31°S, 137°E	rocket	26, 27
White Sands, N. M.	32°N, 106°W	rocket	4, 8, 11, 12, 28-32
Yuma, Ariz.	33°N, 114°W	chemical trails	15, 33-36
Pt. Mugu, Calif.	34°N, 119°W	rocket	5, 37
Adelaide, Australia	35°S, 139°E	meteor radar	38-50

## APPENDIX (continued)

Station	Location	Technique	Reference
Washington, D. C.	37°N, 77°W	I.D.	51, 52
Ashabad, U.S.S.R.	37°N, 58°E	I.D.	53
Dushanbe, U.S.S.R.	37°N, 67°E	meteor radar	54
Arenosillo, Spain	37°N, 6°W	chaff	55-58
Tonopah Range, Nev.	38°N, 117°W	chaff	59
Wallops Island, Va.	38°N, 75°W	rocket, chemical trails	3, 5, 11, 13, 37, 60-71
Chiquita, Argentina	38°S, 57°W	rocket	5
Stamford, Calif.	38°N, 122°W	meteor radar	72
College Park, Pa.	40°N, 77°W	low frequency I.D.	73, 74
Sardinia, Italy	40°N, 90°E	chemical trails	71
Bedford, Mass.	42°N, 71°W	meteor radar	75, 76
Frunze, U.S.S.R.	42°N, 69°E	meteor radar	54, 77
Durham, N. H.	43°N, 70°W	meteor radar	72
Millstone Hill, Mass.	43°N, 71°W	incoherent scatter	78-80
Birdlings Flat, New Zealand	44°S, 173°E	partial reflection I.D.	81
St. Santin, France	45°N, 2°E	incoherent scatter	82-90
Ottawa, Canada	46°N, 76°W	I.D.	21, 52, 91
Garchy, France	47°N, 3°E	meteor radar, I.D.	82, 92-97
Rostov-Don, U.S.S.R.	47°N, 39°E	I.D.	53, 98
Freiburg, Switzerland	47°N, 7°E	I.D.	21, 24, 91, 99, 100
Volgograd, U.S.S.R.	49°N, 45°E	rocket, chemical trails	101, 102
Domont, France	49°N, 2°E	I.D.	91
Kharhov, U.S.S.R.	50°N, 33°E	meteor radar	53, 103
Collin, G. D. R.	51°N, 13°E	low frequency I.D.	104-110
Cambridge, Gr. Br.	52°N, 0°	I.D.	52, 111-113
Swansea, Gr. Br.	52°N, 4°W	I.D.	112, 114, 115
Saskatoon, Canada	52°N, 107°W	partial reflection I.D.	116-118
Debilt, Neth.	52°N, 5°E	I.D.	22, 119-121
Irkutsk, U.S.S.R.	52°N, 104°E	I.D.	53, 122-124
Sheffield, Jodrell Bank	53°N, 2°W	meteor radar	41, 42, 45, 48, 75, 125-128
Kuhlungsborn, G. D. R.	54°N, 12°E	low frequency	104-110
Primrose Lake, Canada	50°N, 110°W	rocket	5
Obninsk, U.S.S.R.	55°N, 35°E	meteor radar	129
Models or data compilations from latitudes 25° to 55°			11, 13, 45, 46, 53, 54, 67, 101, 114, 128, 130-154

TABLE A2. References for Table A1 citations.

1. Rofe, B. <i>et al.</i> , 1967	26. Groves, G., 1965
2. Rahmatullah, M., and S. Jafir, 1971	27. Groves, G., 1967a
3. Reed, R. <i>et al.</i> , 1966a	28. Webb, W., 1964
4. Reed, R., 1967	29. Beyers, N., and B. Miers, 1965
5. World Data Center A, 1969, 1973	30. Reed, R. <i>et al.</i> , 1966b
6. Edwards, H. <i>et al.</i> , 1963a	31. Beyers, N. <i>et al.</i> , 1966
7. Edwards, H. <i>et al.</i> , 1963b	32. Ballard, H. N. <i>et al.</i> , 1972
8. Lenhard, R., and J. Wright, 1963	33. Fuller, R., 1966
9. Rosenberg, N., and H. Edwards, 1964	34. Anthony, M., and W. Epler, 1967
10. Rosenberg, N. <i>et al.</i> , 1964	35. Murphy, C., and G. Bull, 1968
11. Kochanski, A., 1964	36. Wright, J., 1968
12. Miers, B., 1965	37. Mitchell, L., 1970
13. Kochanski, A., 1966	38. Elford, W., and D. Robertson, 1953
14. Rosenberg N., and C. Justus, 1966	39. Elford, W., 1959a
15. Kantor, D., 1968	40. Elford, W., 1959b
16. Woodrum, P., and C. Justus, 1968	41. Haurwitz, O., 1963
17. Rees, D., 1968	42. Haurwitz, O., 1964
18. Woodrum, A. <i>et al.</i> , 1969	43. Roper, R., and W. Elford, 1965
19. Bowen, P. J. <i>et al.</i> , 1964	44. Roper, R., 1966
20. Tsukamoto, K., and Y. Ogata, 1959	45. Groves, G., 1969b
21. Kusherovskiy, Y., and S. Mirhoten, 1961	46. Groves, G., 1971
22. Rao, G., and B. Rao, 1965	47. Lloyd, K., 1972
23. Hurahata, M. <i>et al.</i> , 1966	48. Kochanski, A., 1972
24. Kazimirovskiy, E., 1969	49. Groves, G., 1972
25. Rao, G., and M. Rao, 1972	50. Stubbs, T., 1973

TABLE A2 (continued)

- |   |   |
|---|---|
| 51. Salzberg, C., and D. Greenstone, 1951   | 103. Kascheyev, B., and I. Lysenko, 1968      |
| 52. Briggs, B., and M. Spencer, 1954        | 104. Sprenger, K., 1965                       |
| 53. Mirkoten, S., 1965                      | 105. Sprenger, K., and R. Schminder, 1967     |
| 54. Lysenko, I. <i>et al.</i> , 1969        | 106. Sprenger, K., and R. Schminder, 1968     |
| 55. Azcarraga, A. <i>et al.</i> , 1970      | 107. Sprenger, K. <i>et al.</i> , 1969        |
| 56. Azcarraga, A. <i>et al.</i> , 1971      | 108. Sprenger, K. <i>et al.</i> , 1971        |
| 57. Rose, G. <i>et al.</i> , 1972a          | 109. Sprenger, K., and I. Lysenko, 1972       |
| 58. Rose, G. <i>et al.</i> , 1972b          | 110. Lysenko, I. <i>et al.</i> , 1972         |
| 59. Smith, L. 1960, 1962                    | 111. Phillips, G., 1952                       |
| 60. aufm Kampe, H. S. <i>et al.</i> , 1962  | 112. Beynon, W., and G. Goodwin, 1958         |
| 61. Hines, C., 1966                         | 113. Jones, I., 1958                          |
| 62. Finger, P., and H. Woolf, 1967          | 114. Goodwin, G., 1968                        |
| 63. Theon, J. <i>et al.</i> , 1966          | 115. Beynon, W. <i>et al.</i> , 1972          |
| 64. Bedinger, J., and L. Smith, 1967        | 116. Gregory, J., and D. Rees, 1971           |
| 65. Bedinger, J. <i>et al.</i> , 1968       | 117. Manson, A. <i>et al.</i> , 1973          |
| 66. Tarpley, J., 1970                       | 118. Manson, A. <i>et al.</i> , 1974          |
| 67. Fnttsian, G., and D. Tarasenko, 1971    | 119. Harang, L., 1964                         |
| 68. Cole, A., and A. Kantor, 1972           | 120. Harang, L., 1965                         |
| 69. Theon, J. <i>et al.</i> , 1972          | 121. Vesseur, H., 1972                        |
| 70. Hoxit, L., and R. Henry, 1973           | 122. Kazimirovskiy, E., and V. Kokourov, 1961 |
| 71. Kochanski, A., 1973                     | 123. Kokourov, V. <i>et al.</i> , 1971        |
| 72. Barnes, A., 1972                        | 124. Kazimirovskiy, E., and V. Kokourov, 1972 |
| 73. Sales, G., and S. Bowhill, 1962         | 125. Greenhow, J., and E. Newfeld, 1956       |
| 74. Lee, II., and S. Bowhill, 1963          | 126. Greenhow, J., and E. Neufeld, 1961       |
| 75. Barnes A., and J. Pazniokas, 1972       | 127. Müller, H., 1966                         |
| 76. Barnes A., 1973                         | 128. Kent, G., and R. Wright, 1968            |
| 77. Lysenko, I. <i>et al.</i> , 1972        | 129. Iljichev, Y. <i>et al.</i> , 1974        |
| 78. Salah, J. E. <i>et al.</i> , 1974       | 130. Batten, E., 1961                         |
| 79. Salah, J. E., 1974                      | 131. Kazimirovskiy, E., 1962                  |
| 80. Salah, J. E. <i>et al.</i> , 1975       | 132. Kazimirovskiy, E., 1963                  |
| 81. Fraser, G., and A. Kochanski, 1970      | 133. Kantor, A., and A., Cole 1964            |
| 82. Carru, H. <i>et al.</i> , 1967          | 134. Kantor, A., 1965                         |
| 83. Waldteufel, P., 1970                    | 135. Rawer, K., 1965b                         |
| 84. Bernard, R., and A. Spizzichino, 1971   | 136. Henderson, L., 1965                      |
| 85. Waldteufel, P. <i>et al.</i> , 1972     | 137. Bhatnagar, V., and P. Mitra, 1966        |
| 86. Bernard, R., 1974a                      | 138. Anderson, A., and W. Francis, 1966       |
| 87. Amayenc, P., and C. Reddy, 1974         | 139. Champion, K., 1967                       |
| 88. Amayenc, P., 1974                       | 140. Groves, G., 1967b                        |
| 89. Fontani, J., and D. Alcayde, 1974       | 141. Reddy, C., and S. Matsushita, 1968       |
| 90. Bernard, R., 1974b                      | 142. Rosenberg, N., 1968                      |
| 91. Rawer, K., 1965a                        | 143. Groves, G., and S. Makarious, 1968       |
| 92. Spizzichino, A., 1967                   | 144. Rao, G., and B. Rao, 1964                |
| 93. Spizzichino, A., and J. Revah, 1968     | 145. Reed, R. <i>et al.</i> , 1969            |
| 94. Spizzichino, A., 1969                   | 146. Groves, G., 1969a                        |
| 95. Revah, I., 1969                         | 147. Chapman, S., and R. Lindzen, 1970        |
| 96. Haubert, A. <i>et al.</i> , 1971        | 148. Groves, G., 1970a                        |
| 97. Fellous, J. <i>et al.</i> , 1974        | 149. Groves, G., 1970b                        |
| 98. Pozigun, V., 1968                       | 150. Jacchia, L., 1971                        |
| 99. Harnischmacher, E., 1964                | 151. Reed, A., 1972                           |
| 100. Harnischmacher, E., and K. Rawer, 1968 | 152. Teptin, G., 1972                         |
| 101. Bugayeva, I., and L. Ryazanova, 1971   | 153. Rosenberg, N., and S. Zimmerman, 1972    |
| 102. Andreeva, L. <i>et al.</i> , 1972      | 154. MacDougall, J., 1974                     |

## REFERENCES

- Amayenc, P., 1974: Tidal oscillations of the meridional neutral wind at mid latitudes. *Radio Sci.*, **9**, 281-291.
- , and C. A. Reddy, 1972: Height structure of tidal winds as inferred from incoherent scatter observations. *Planet. Space Sci.*, **20**, 1269-1279.
- Anderson, A., and W. Francis, 1966: The variation of the neutral atmospheric properties with local time and solar activity from 100 to 10,000 km. *J. Atmos. Sci.*, **23**, 110-124.
- Andreeva, L. A. *et al.*, 1972: Wind profiles from artificial luminous cloud observations and the theory of  $E_s$  formation based on wind shear. *Phil. Trans. Roy. Soc. London*, **A271**, 623-629.
- Anthony, M., and W. Epler, 1967: Upper atmosphere winds from gun launched vertical probes (Yuma 16-19 Nov. 1966). Space Instruments Research Inc., Atlanta, Ga. [Available from DDC, AD 662719, 85 pp.]
- aufm Kampe, H., M. Smith and R. Brown, 1962: Winds between 60 and 110 km. *J. Geophys. Res.*, **67**, 4243-4258.
- Azcarraga, A., L. Sanchez and H. Widdel, 1970: On the structure of mesospheric winds. *Space Research X*, North-Holland, 174-179.
- , — and —, 1971: Measured wind oscillation at mesospheric levels. *Space Research XI*, North-Holland, 835-840.
- Ballard, H. *et al.*, 1972: Atmospheric tidal measurements at 50 km from a constant-altitude balloon. *J. Appl. Meteor.*, **11**, 1138-1149.
- Barnes, A. A. Jr., 1972: Radar meteor trail task—Final report. AFCRL-72-0190, Environ. Res. Pap., No. 397, 46 pp.
- , 1973: Status report on radar meteor wind and density measurements. *Bull. Amer. Meteor. Soc.*, **54**, 900-911.

- , and J. Pazniokas, 1972: Results from the AFCRL radar meteor trail set. AFCRL-72-0185, Environ. Res. Pap., No. 392, 81 pp.
- Bartten, E., 1961: Wind systems in the mesosphere and lower ionosphere. *J. Meteor.*, **18**, 283-291.
- Bedinger, J., and L. Smith, 1967: Measured wind shears and electron densities. *Space Research VII*, North-Holland, 100-112.
- , et al., 1968: Upper atmosphere winds and their interpretation-I. Evidence for strong nonlinearity of the horizontal flow above 80 km. *Planet. Space Sci.*, **16**, 159-193.
- Bernard, R., 1974a: Tides in the E region observed by incoherent scatter over St. Santin. *J. Atmos. Terr. Phys.*, **36**, 1105-1120.
- , 1974b: A comparison between meteoric radar and incoherent scatter measurements in the lower thermosphere. *Radio Sci.*, **9**, 295-300.
- , and A. Spizzichino, 1971: Semi-diurnal wind and temperature oscillations in the E-region observed by the Nancy incoherent scatter equipment. *J. Atmos. Terr. Phys.*, **33**, 1345-1352.
- Byers, N., and B. Miers, 1965: Diurnal temperature change in the atmosphere between 30 and 60 km over White Sands Missile Range. *J. Atmos. Sci.*, **22**, 262-266.
- , and R. Reed, 1966: Diurnal tidal motions near the stratopause during 48 hours at White Sands Missile Range. *J. Atmos. Sci.*, **23**, 325-333.
- Beynon, W., and G. Goodwin, 1958: Horizontal drifts and temperature in the lower part of region E. *J. Atmos. Terr. Phys.*, **13**, 180-182.
- , M. Barrett and R. Edean, 1972: Some studies of sporadic E-layer drifts. *Phil. Trans. Roy. Soc. London*, **A271**, 613-622.
- Bhatnagar, V., and A. Mitra, 1966: An upper atmosphere model for solar minimum conditions. *J. Atmos. Sci.*, **23**, 233-244.
- Bowen, P. et al., 1964: Rocket studies of sporadic-E ionization and ionospheric winds. *Planet. Space Sci.*, **12**, 1173-1177.
- Briggs, B., and M. Spencer, 1954: Horizontal movements in the ionosphere. *Rep. Prog. Phys.*, **17**, 245-265.
- Bugayeva, I., and L. Ryazanova, 1971: Annual and diurnal variability of the temperature field in the 30-80 km layer. *Meteor. Gidrol.*, **12**, 44-53.
- Carru, H. et al., 1967: Resultats ionospheriques obtenus par diffusion de Thomson (1966-1967). *Ann. Geophys.*, **23**, 455-466.
- Champion, K., 1967: Variations with season and latitude of density, temperature and composition in the lower thermosphere. *Space Research VII*, North-Holland, 1101-1118.
- Chapman, S., and R. Lindzen, 1970: *Atmospheric tides*. Gordon and Breach, 200 pp.
- Cole, A., and A. Kantor, 1972: COESA task group I report on proposed 50 to 100 km revision to U. S. Standard Atmosphere 1962. AFCRL-72-0692, ERP, 422.
- Donn, W. L., and D. Rind, 1972: Microbaroms and the temperature and wind of the upper atmosphere. *J. Atmos. Sci.*, **29**, 156-172.
- , and B. Naini, 1973: Sea wave origin of microbaroms and microseisms. *J. Geophys. Res.*, **78**, 4482-4488.
- Edwards, H. et al., 1963: Upper atmosphere wind measurements determined from twelve rocket experiments. *J. Geophys. Res.*, **68**, 3021-3032.
- , Justus, C. and D. Kurts, 1963b: Evening twilight winds from 68 to 140 km for May 21, 1963. *J. Geophys. Res.*, **68**, 6062-6063.
- Elford, W., 1959a: A study of winds between 80 and 100 km in medium latitudes. *Planet. Space Sci.*, **1**, 94-101.
- , 1959b: Winds in the upper atmosphere. *J. Atmos. Terr. Phys.*, **15**, 132-136.
- , and D. Robertson, 1953: Measurements of winds in the upper atmosphere by means of drifting meteor trails II. *J. Atmos. Terr. Phys.*, **4**, 271-284.
- Entsian, G., and D. Tarasenko, 1971: Seasonal rearrangement of circulation and processes in the ionosphere. *Meteor. Gidrol.*, **5**, 23-32.
- Fellous, J. et al., 1974: Vertical propagation of tides at meteor heights. *J. Atmos. Terr. Phys.*, **36**, 385-397.
- Finger, F., and H. Woolf, 1967: Diurnal variation of temperature in the upper stratosphere as indicated by a meteorological rocket experiment. *J. Atmos. Sci.*, **24**, 230-239.
- Fontanari, J., and D. Alcayde, 1974: Observation of neutral temperature tidal-type oscillations in the F<sub>1</sub> region. *Radio Sci.*, **9**, 275-280.
- Fraser, G., and A. Kochanski, 1970: Ionospheric drifts from 64-108 km altitudes at Birdlings Flat. *Ann. Geophys.*, **26**, 675-687.
- Fuller, R., 1966: Upper atmosphere winds from gun launched vertical probes (Yuma 13-15 6/66). Space Instruments Research, Inc. Atlanta, Ga. [Available from DDC, AD 662 717, 38 pp.].
- Goodwin, G., 1968: Fourier analysis and interpretation of D- and E- region winds. *J. Atmos. Terr. Phys.*, **30**, 995-1018.
- Greenhow, J., and E. Neufeld, 1956: The height variation of upper atmospheric winds. *Phil. Mag.*, **1**, 1157-1171.
- , and —, 1961: Winds in the upper atmosphere. *Quart. J. Roy. Meteor. Soc.*, **87**, 472-489.
- Gregory, J., and D. Rees, 1971: Wind Profile to 100 km near 53°N during 1969. *J. Atmos. Sci.*, **28**, 1079-1083.
- Groves, G., 1965: Diurnal variations in upper atmosphere wind and temperature structure at Woomera, 15-16 October 1963. *Space Research V*, North-Holland 1012-1020.
- , 1967a: Variations in upper atmospheric wind, temperature and pressure at Woomera during the night of 29/30 April 1965. *Space Research VII*, North-Holland, 977-985.
- , 1967b: Diurnal and semidiurnal S-N wind components in the upper atmosphere derived from rocket measurements. *Space Research VII*, North-Holland, 986-1000.
- , 1969a: Wind models from 60-130 km altitude for different months and latitudes. *J. Brit. Interplanet. Soc.*, **22**, 285-307.
- , 1969b: Tidal oscillations and gravity waves. *Ann. IQSY*, **5**, 263-289.
- , 1970a: Seasonal and latitudinal models of atmospheric temperature, pressure and density, 25-110 km. AFCRL-70-0261, AF Survey Geophys. No. 218, 70 pp.
- , 1970b: Seasonal and latitudinal models of atmospheric structure between 30 and 120 km. *Space Research X*, North-Holland 137-150.
- , 1971: Atmospheric structure and its variations in the region from 25 to 120 km. AFCRL-71-0410, ERP 368, 200 pp.
- , 1972: Annual and semi-annual zonal wind components and corresponding temperature and density variations, 60-130 km. *Planet. Space Sci.*, **20**, 2099-2112.
- , and S. Makarios, 1968: Diurnal S-N wind components below 60 km derived from rocket observations for various seasons and latitudes. *Space Research VIII*, North-Holland, 857-864.
- Harang, L., 1964: Drift measurements of the E and E<sub>s</sub> at high latitudes. *Ann. Geophys.*, **20**, 435-446.
- , 1965: Drift measurements of the E-region in Norway at Kjeller and Tromsø. *Ann. IGY*, **33**, 271-278.
- Harnischmacher, E., 1964: Evolution dans les temps des vents dans la region E. *Ann. Geophys.*, **20**, 425-433.
- , and K. Rawer, 1968: Some observed phenomenon to be explained by theories of ionospheric drift. *J. Atmos. Terr. Phys.*, **30**, 871-884.
- Haubert, A., J. Tabbagh and G. Doyen, 1971: Les vents ionospheriques de la region E. Observations faits a Garchy, par la methode des fadings, en 1969 et 1970. Tech. Note GRI/NTP/87.
- Haurwitz, B., 1963: Wind and pressure oscillations in the upper atmosphere. *Arch. Meteor. Geophys. Bioklim.*, **A13**, 144-166.
- , 1964: Tidal phenomena in the upper atmosphere. WMO Tech Note 58, 27 pp.

- Henderson, L., 1965: Seasonal and latitude variations of noon bearings of E-region drifts. *Ann. IGY*, **33**, 289-292.
- Hines, C., 1966: Diurnal tide in the upper atmosphere. *J. Geophys. Res.*, **71**, 1453-1459.
- Hoxit, L., and R. Henry, 1973: Diurnal and annual temperature variations in the 30-60 km region as indicated by statistical analysis of rocket and temperature data. *J. Atmos. Sci.*, **30**, 922-933.
- Huruhata, M., J. Nakamura and K. Saito, 1966: Measurements of upper atmospheric wind by sodium cloud drifts. *Rep. Ionos. Space Res. Jap.*, **20**, 214-222.
- Ilijchev, Y. et al., 1974: Space and time variations of large-scale air motions in the lower thermosphere. *J. Atmos. Terr. Phys.*, **36**, 1841-1850.
- Jacchia, L. G., 1971: Revised static models of the thermosphere and exosphere with empirical temperature profiles. *Smith Astrophys. Obs. Spec. Rep.*, No. 332, 113 pp.
- Jones, I. L., 1958: The height variation of drift in the E-region. *J. Atmos. Terr. Phys.*, **12**, 68-76.
- Kantor, A., 1965: Interim notes on atmospheric properties.—Solar effects on winds 80-130 km an interim analysis. AFCRL INAP 49, 25 pp.
- , 1968: Winds above 85 km, Eglin AFB, Fla. and Yuma, Arizona. AFCRL, INAP 85, 10 pp.
- , and A. Cole, 1964: Zonal and meridional winds to 120 km. *J. Geophys. Res.*, **69**, 5131-5140.
- Kashcheyev, B., and I. Lysenko, 1968: Atmospheric circulation in the meteor zone. *J. Atmos. Terr. Phys.*, **30**, 903-906.
- Kazimirovskiy, E., 1962: The problem of general circulation in the ionosphere. *Geomagn. Aeronomiya*, **2**, 6, 897-902.
- , 1963: Wind systems in the lower ionosphere. *Geomagn. Aeronomiya*, **3**, 380-387.
- , 1969: Horizontal ionospheric drifts during low solar activity. *Geomagn. Aeronomiya*, **9**, 839-845.
- , and V. Kokourov, 1961: Research on the irregular structure of the ionosphere above Irkutsk during the IGY. *Geomagn. Aeronomiya*, **1**, 654-662.
- , and —, 1972: The measurement of ionospheric drifts over East Siberia, USSR. *Phil. Trans. Roy. Soc. London*, **A271**, 499-508.
- Kent, G., and R. Wright, 1968: Movements of ionospheric irregularities and atmospheric winds. *J. Atmos. Terr. Phys.*, **30**, 657-692.
- Kochanski, A., 1964: Atmospheric motions from sodium cloud drifts. *J. Geophys. Res.*, **69**, 3651-3662.
- , 1966: Atmospheric motions from sodium cloud drifts at four locations. *Mon. Wea. Rev.*, **94**, 199-212.
- , 1972: Semiannual variation at the base of the thermosphere. *Mon. Wea. Rev.*, **100**, 222-234.
- , 1973: Tidal motions in the ionosphere from sodium cloud drifts. *Ann. Geophys.*, **29**, 77-87.
- Kokourov, V. et al., 1971: The simultaneous measurements of ionospheric drifts by vertical and oblique sounding. *J. Atmos. Terr. Phys.*, **33**, 943-950.
- Kushnerovskiy, Y., and S. Mirkotan, 1961: Motions of ionospheric irregularities: a survey. *Geomagn. Aeronomiya*, **1**, 405-426.
- Lee, H., and S. Bowhill, 1963: A statistical study of apparent horizontal ionospheric movements using 300 kc/s radio waves. *J. Atmos. Terr. Phys.*, **25**, 659-685.
- Lenhard, R., and J. Wright, 1963: Mesospheric winds from 23 successive hourly soundings. *J. Geophys. Res.*, **68**, 227-234.
- Lloyd, K. H., 1972: Observed combined chemical seeding and ground-based technique—I. Winds, turbulence and parameters of the neutral atmosphere. *Planet. Space Sci.*, **20**, 761-789.
- Lysenko, I. et al., 1969: Results of radar-meteor tracking studies on atmospheric circulation over Eurasia and the Arctic. *Izv. Atmos. Oceanic Phys.*, **5**, 508-513.
- , A. Orlyansky and Y. Portnyagin, 1972: A study of the wind regime at an altitude of about 100 km by the meteor-radar method. *Phil. Trans. Roy. Soc. London*, **A271**, 601-612.
- Lysenko, I. et al., 1972: Results of a comparison between radar meteor wind measurements and simultaneous lower ionosphere drift measurements in the same area. *J. Atmos. Terr. Phys.*, **34**, 1435-1445.
- MacDougall, J., 1974: 110 km neutral zonal wind patterns. *Planet. Space Sci.*, **22**, 545-558.
- Manson, A., J. Gregory and D. Stephenson, 1973: Winds and wave motions (90-100 km) as measured by a partial reflection radiowave system. *J. Atmos. Terr. Phys.*, **35**, 2055-2078.
- , — and —, 1974: Winds and wave motions to 110 km at mid-latitudes. I. Partial reflection radiowave soundings, 1972-73. *J. Atmos. Sci.*, **31**, 2207-2215.
- Miers, B., 1965: Wind oscillations between 30 and 60 km over White Sands Missile Range, New Mexico. *J. Atmos. Sci.*, **22**, 382-387.
- Mirkotan, S., 1965: The main results of investigation of the ionospheric drifts in the USSR for the IGY/IGC. *Ann. IGY*, **33**, 243-264.
- Mitchell, L., 1970: Variability of the monthly mean zonal wind, 30-60 km. AWS Tech. Rept. 195, 80 pp.
- Müller, H., 1966: Atmospheric tides in the meteor zone. *Planet. Space Sci.*, **14**, 1253-1272.
- Murphy, C., and G. Bull, 1968: Ionospheric winds over Yuma, Arizona, measured by gun-launched projectiles. *J. Geophys. Res.*, **73**, 3005-3015.
- Phillips, G., 1952: Measurement of winds in the ionosphere. *J. Atmos. Terr. Phys.*, **2**, 141-154.
- Posmentier, E., 1967: A theory of microbaroms. *Geophys. J. Roy. Astron. Soc.*, **13**, 487-501.
- Pozigun, V., 1968: Description of drifts and of the inhomogeneous structure of the ionosphere over Rostov-en-the-Don during the IGY-IQSY period. *Geomagn. Aeronomiya*, **8**, 365-368.
- Rahmatullah, M., and S. Jafri, 1971: Mean stratospheric wind circulation at Sonmiani and some peculiarities observed during 1969-1970. *Space Research XI*, North-Holland, 807-812.
- Rao, G., and B. Rao, 1964: The latitude variation of height gradient in the E-region drift speeds. *J. Atmos. Terr. Phys.*, **26**, 628-629.
- , and —, 1965: The latitude variation of apparent horizontal movements in the E region of the ionosphere. *J. Geophys. Res.*, **70**, 667-677.
- , and —, 1972: Horizontal drifts in E- and E<sub>s</sub>-layers for middle latitude station. *J. Atmos. Terr. Phys.*, **34**, 537-541.
- Rawer, K., 1965a: (Compiler) Results of ionospheric drift observations. *Ann. IGY*, **33**, 1-242.
- , 1965b: Intercomparison of instantaneous drift data. *Ann. IGY*, **33**, 293-296.
- Reddy, C., and S. Matsushita, 1968: The variations of neutral wind shears in the E-region as deduced from blanketing E<sub>s</sub>. *J. Atmos. Terr. Phys.*, **30**, 747-762.
- Reed, R., 1967: Semidiurnal tidal motions between 30 and 60 km. *J. Atmos. Sci.*, **24**, 315-317.
- , 1972: Further analysis of semidiurnal tidal motions between 30 and 60 km. *Mon. Wea. Rev.*, **100**, 579-581.
- , J. McKenzie and J. Vyverberg, 1966a: Diurnal tidal motions between 30 and 60 km in summer. *J. Atmos. Sci.*, **23**, 416-423.
- , — and —, 1966b: Further evidence of enhanced diurnal tidal motions near the stratopause. *J. Atmos. Sci.*, **23**, 247-251.
- , M. Oard and M. Sieminski, 1969: A comparison of observed and theoretical diurnal tidal motions between 30 and 60 km. *Mon. Wea. Rev.*, **97**, 456-459.
- Rees, D., 1968: Determinations of upper atmosphere structure above 90 km by optical observations on rocket-borne chemical releases. *Space Research VIII*, North-Holland, 909-919.
- Revah, I., 1969: Etude des vents de petites échelles observés au moyen des traînées météoriques. *Ann. Geophys.*, **25**, 1-45.

- Rind, D., W. L. Donn and E. Dede, 1973: Upper air wind speeds calculated from observations of natural infrasound. *J. Atmos. Sci.* **30**, 1726-1729.
- Rofe, B., W. Elford and E. Doyle, 1967: Daily variations in winds, density, temperature and pressure between 40 and 90 km, in the sub-tropical latitudes of the Southern Hemisphere. (abstract only). *Space Research VII*, North-Holland, 973-976.
- Roper, R., 1966: The semidiurnal tide in the lower thermosphere. *J. Geophys. Res.*, **71**, 5746-5748.
- , and W. Elford, 1965: Meteor winds measured at Adelaide (35°S) 1961. NASA X-650-65-220.
- Rose, G. et al., 1972a: Results of an experimental investigation of correlations between D region neutral gas winds, density changes, and short wave absorption. *Radio Sci.*, **7**, 181-191.
- , et al., 1972b: A payload for small sounding rockets for wind finding and density measurements in the height region between 95 and 75 km. *Phil. Trans. Roy. Soc. London*, **A271**, 509-528.
- Rosenberg, N., 1968: Ionospheric winds—a statistical analysis. *Space Research VIII*, North-Holland, 673-678.
- , and H. Edwards, 1964: Observations of ionospheric wind patterns through the night. *J. Geophys. Res.*, **69**, 2819-2826.
- , —, and J. Wright, 1964: Ionospheric winds: motions into night and sporadic E correlations. *Space Research IV*, North-Holland, 171-181.
- , and C. Justus, 1966: Space and time correlations of ionospheric winds. *Radio Sci.*, **1**, 149-155.
- , and S. Zimmerman, 1972: Ionospheric winds and vicious dissipation. *Radio Sci.*, **7**, 377-380.
- Salah, J., 1974: Daily oscillations of the mid-latitude thermosphere studied by incoherent scatter at Millstone Hill. *J. Atmos. Terr. Phys.*, **36**, 1891-1910.
- , J. Evans and R. Ward, 1974: Seasonal variations in the thermosphere above Millstone Hill. *Radio Sci.*, **9**, 231-238.
- , —, and —, 1975, E-region temperature measurements at Millstone Hill. *J. Atmos. Terr. Phys.*, **37**, 461-490.
- Sales, G., and S. Bowhill, 1962: Apparent ionospheric motion in the night-time D-region. *J. Atmos. Terr. Phys.*, **24**, 451-465.
- Salzberg, C., and R. Greenstone, 1951: Systematic ionospheric winds. *J. Geophys. Res.*, **56**, 521-533.
- Smith, L., 1960: The measurement of winds between 100,000 and 300,000 feet by use of chaff rockets. *J. Meteor.*, **17**, 296-310.
- , 1962: Monthly wind measurements in the mesodecline over a one-year period. *J. Geophys. Res.*, **67**, 4653-4672.
- Spizzichino, A., 1967: Mesures de vents ionosphériques au moyen d'un radar météorologique. *Space Research VII*, North-Holland, 73-75.
- , 1969: Etude expérimentales des vents dans la haute atmosphère. Part I. *Ann. Geophys.*, **25**, 697-720.
- , and I. Revah, 1968: Nouvelles données expérimentales sur les différentes composantes du vent au moyen d'un radar météorologique et discussion des résultats obtenus au moyen de fusées. *Space Research VIII*, North-Holland, 679-682.
- Sprenger, K., 1965: Seasonal anomalies of drift directions in the lower ionosphere revealed by spaced receiver measurements in the LF range. *Ann. IGY*, **33**, 266-270.
- , and R. Schindler, 1967: Results of ten years' ionospheric drift measurements in the l.f. range. *J. Atmos. Terr. Phys.*, **29**, 183-199.
- , and —, 1968: On the significance of ionospheric drift measurements in the LF range. *J. Atmos. Terr. Phys.*, **30**, 693-700.
- , K. Greisiger and R. Schindler, 1969: Seasonal changes in phase of the semidiurnal tidal wind component in the lower ionosphere. *Ann. Geophys.*, **25**, 505-509.
- , —, and —, 1971: Seasonal variations in the semi-diurnal tidal wind determined from observations of drift of ionospheric inhomogeneities at altitude of 90-110 km. *Izv. Atmos. Oceanic Phys.*, **7**, 311-316.
- , and I. Lysenko, 1972: The significance of ionospheric drift measurements in the low-frequency range. *Phil. Trans. Roy. Soc. London*, **A271**, 473-484.
- Stubbs, T., 1973: The measurement of winds in the D-region of the ionosphere by the use of partially reflected radio waves. *J. Atmos. Terr. Phys.*, **35**, 909-919.
- Tarpley, J., 1970: The ionospheric wind dynamo-II solar tides. *Planet. Space Sci.*, **18**, 1091-1103.
- Teptin, G., 1972: Upper atmosphere motion, as determined by observations of radio echos from meteor trails. *Izv. Atmos. Oceanic Phys.*, **8**, 139-150.
- Theon, J., 1972: The mean observed meteorological structure and circulation of the stratosphere and mesosphere. NASA TR R-375, 77 pp.
- , et al., 1966: Some observations on the thermal behavior of the mesosphere. NASA X-621-66-490, 36 pp.
- Tsukamoto, K., and Y. Ogata, 1959: Ionospheric drifts at Yamagawa in Japan. *Rep. Ionos. Space Res. Jap.*, **13**, 48-55.
- Vesseyer, H., 1972: Results of E-layer drift measurements at De Bilt. *Phil. Trans. Roy. Soc. London*, **A271**, 485-488.
- Waldfteufel, P., 1970: A study of seasonal changes in the lower thermosphere and their implications. *Planet. Space Sci.*, **18**, 741-748.
- , P. Bauer and J. McClure, 1972: Structure of the thermosphere as inferred from incoherent scatter measurements. *Space Research XII*, North-Holland, 899-907.
- Webb, W., 1964: Stratospheric solar response. *J. Atmos. Sci.*, **21**, 582-591.
- Woodrum, A., and C. Justus, 1968: Atmospheric tides in the height region 90-120 km. *J. Geophys. Res.*, **73**, 467-478.
- , —, and R. Roper, 1969: Measurements of tidal-period winds in the 95-135 km region. *J. Geophys. Res.*, **74**, 4099-4104.
- World Data Center A—Meteorology, 1969, 1973: High Altitude Meteorological Data. U. S. Department of Commerce, NOAA.
- Wright, J., 1968: The interpretation of ionospheric radio drift measurements-I. Some results of experimental comparisons with neutral wind profiles. *J. Atmos. Terr. Phys.*, **30**, 919-930.